

**ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Қ. И. Сәтпаев атындағы Қазақ ұлттық техникалық зерттеу университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Казахский национальный исследовательский
технический университет им. К. И. Сатпаева

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Kazakh national research technical university
named after K. I. Satpayev

SERIES
OF GEOLOGY AND TECHNICAL SCIENCES

6 (438)

NOVEMBER – DECEMBER 2019

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруды. Web of Science зерттеушілер, авторлар, баспашилар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Emerging Sources Citation Index-ке енүі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

Бас редакторы
э. ф. д., профессор, КР ҮГА академигі
И.К. Бейсембетов
Бас редакторының орынбасары
Жолтаев Г.Ж. проф., геол.-мин. ф. докторы
Редакция ақысы:

Абаканов Т.Д. проф. (Қазақстан)
Абишева З.С. проф., академик (Қазақстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Әзірбайжан)
Бакиров А.Б. проф., (Қыргыстан)
Беспаев Х.А. проф. (Қазақстан)
Бишимбаев В.К. проф., академик (Қазақстан)
Буктуков Н.С. проф., академик (Қазақстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Тәжікстан)
Грэвис Р.М. проф. (АҚШ)
Ерғалиев Г.К. проф., академик (Қазақстан)
Жуков Н.М. проф. (Қазақстан)
Қожахметов С.М. проф., академик (Казахстан)
Конторович А.Э. проф., академик (Ресей)
Курскеев А.К. проф., академик (Қазақстан)
Курчавов А.М. проф., (Ресей)
Медеу А.Р. проф., академик (Қазақстан)
Мұхамеджанов М.А. проф., корр.-мүшесі (Қазақстан)
Нигматова С.А. проф. (Қазақстан)
Оздоев С.М. проф., академик (Қазақстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Қазақстан)
Сейтов Н.С. проф., корр.-мүшесі (Қазақстан)
Сейтмуратова Э.Ю. проф., корр.-мүшесі (Қазақстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (АҚШ)
Штейнер М. проф. (Германия)

«ҚР ҮГА Хабарлары. Геология мен техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РКБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрагат комитетінде 30.04.2010 ж. берілген №10892-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://www.geolog-technical.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыра көш., 69а.

мекенжайы: Қ. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «Аруна» ЖҚ, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д. э. н., профессор, академик НАН РК

И. К. Бейсембетов

Заместитель главного редактора

Жолтаев Г.Ж. проф., доктор геол.-мин. наук

Р е д а к ц и о н а я к о л л е г и я:

Абаканов Т.Д. проф. (Казахстан)
Абишева З.С. проф., академик (Казахстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Азербайджан)
Бакиров А.Б. проф., (Кыргызстан)
Беспаев Х.А. проф. (Казахстан)
Бишимбаев В.К. проф., академик (Казахстан)
Буктуков Н.С. проф., академик (Казахстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Таджикистан)
Грэвис Р.М. проф. (США)
Ергалиев Г.К. проф., академик (Казахстан)
Жуков Н.М. проф. (Казахстан)
Кожахметов С.М. проф., академик (Казахстан)
Конторович А.Э. проф., академик (Россия)
Курскеев А.К. проф., академик (Казахстан)
Курчавов А.М. проф., (Россия)
Медеу А.Р. проф., академик (Казахстан)
Мухамеджанов М.А. проф., чл.-корр. (Казахстан)
Нигматова С.А. проф. (Казахстан)
Оздоев С.М. проф., академик (Казахстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Казахстан)
Сеитов Н.С. проф., чл.-корр. (Казахстан)
Сейтмуратова Э.Ю. проф., чл.-корр. (Казахстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (США)
Штейнер М. проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: Республикаинское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10892-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© Национальная академия наук Республики Казахстан, 2019

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief
doctor of Economics, professor, academician of NAS RK

I. K. Beisembetov

Deputy editor in chief

Zholtayev G.Zh. prof., dr. geol-min. sc.

Editorial board:

Abakanov T.D. prof. (Kazakhstan)
Abisheva Z.S. prof., academician (Kazakhstan)
Agabekov V.Ye. academician (Belarus)
Aliyev T. prof., academician (Azerbaijan)
Bakirov A.B. prof., (Kyrgyzstan)
Bespayev Kh.A. prof. (Kazakhstan)
Bishimbayev V.K. prof., academician (Kazakhstan)
Buktukov N.S. prof., academician (Kazakhstan)
Bulat A.F. prof., academician (Ukraine)
Ganiyev I.N. prof., academician (Tadzhikistan)
Gravis R.M. prof. (USA)
Yergaliев G.K. prof., academician (Kazakhstan)
Zhukov N.M. prof. (Kazakhstan)
Kozhakhmetov S.M. prof., academician (Kazakhstan)
Kontorovich A.Ye. prof., academician (Russia)
Kurskeyev A.K. prof., academician (Kazakhstan)
Kurchavov A.M. prof., (Russia)
Medeu A.R. prof., academician (Kazakhstan)
Muhamedzhanov M.A. prof., corr. member. (Kazakhstan)
Nigmatova S.A. prof. (Kazakhstan)
Ozdoyev S.M. prof., academician (Kazakhstan)
Postolatii V. prof., academician (Moldova)
Rakishev B.R. prof., academician (Kazakhstan)
Seitov N.S. prof., corr. member. (Kazakhstan)
Seitmuratova Ye.U. prof., corr. member. (Kazakhstan)
Stepanets V.G. prof., (Germany)
Humphery G.D. prof. (USA)
Steiner M. prof. (Germany)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 10892-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-namrk.kz/geology-technical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Editorial address: Institute of Geological Sciences named after K.I. Satpayev
69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 6, Number 438 (2019), 96 – 103

<https://doi.org/10.32014/2019.2518-170X.160>

UDC 551.58:631+551.4

IRSTI 87.29.91

G. A. Adilbektegi¹, J. S. Mustafayev², T. K. Uvatayeva¹, Z. N. Dulatbekova¹, Jozef Mosiej³

¹RSE L. N. Gumilyov Eurasian National University, Astana, Kazakhstan,

²JSC Kazakh National Agrarian University, Almaty, Kazakhstan,

³Warsaw University of Natural Sciences, Warsaw, Poland.

E-mail: adilbekova.gulmira@yandex.kz ,z-mustafa@rambler.ru,
yadulatbekova@gmail.com, t_uvatayeva@mail.ru, jozef_mosiej@sggw.pl

**QUANTITATIVE AND QUALITATIVE ASSESSMENT
OF BIOLOGICAL AND ECOLOGICAL POTENTIAL
OF THE LANDSCAPES OF SOUTHERN KAZAKHSTAN**

Abstract. The paper considers methodological approaches to assessment of biological and environmental potential of landscape systems in the southern Kazakhstan soil-climatic conditions to the most complete and efficient use of natural resources in the design and construction of high-performance agrolandscape systems. Based on qualitative and quantitative indicators heat, light and moisture availability a mathematical model has been developed to assess biological and environmental productivity of landscape systems including soil productivity and vegetation cover.

It should be noted that the developed model of landscapes biological productivity assessment (C_{bp}) takes into account the heat availability of the landscape systems soil cover (C_{tb}), natural moisturization coefficient of the vegetation cover (C_m), and ecological productivity of landscapes (C_{bp}). This model is based on the potentially available energy expended in soil-forming process (Q_n) and natural moisturization coefficient of the vegetation cover (C_m) is one of the modifications of the qualitative and quantitative models of natural system climatic productivity, widely used in the field of geography and ecology assessment of natural system resources potential.

Based on the developed models of landscape systems biological and environmental productivity a comprehensive assessment of South Kazakhstan region natural resource potential was performed, including Almaty, Zhambyl, Kyzylorda and Turkestan region using long-term data of 32 meteorological stations located on the territory of the region.

Use of improved methodologies for assessment of landscape systems biological and ecological productivity can more accurately determine the biological resources of the landscape systems according to agro-ecological areas and to assess the efficiency of natural system natural resources potential.

Keywords: nature, landscape, climate, productivity, biology, ecology, potential, index.

Introduction. To solve the problems of rational distribution of agriculture productive forces and designing high-performance agrolandscape systems, it is necessary to have a detailed landscapes descriptions by the most important factors characterizing heat and moisture availability, plant and soil covers, expressed in the form of some mathematical models to evaluate their natural resource potential.

Existing methodological approaches and methods of the climate productivity assessment do not adequately reflect the productivity of landscape systems, i.e., vegetation and soil cover, there is a need to develop methods of complex biological and ecological assessment of the landscapes productivity on the basis of the fundamental laws of nature, which must include a private assessment of its components productivity, that is, soil and vegetative cover productivity.

Biological and environmental assessment productivity landscapes should be understood as a comprehensive assessment using the integral characteristics of climate, soil and environmental factors positively influencing the growth and development of plants in certain geographic areas representing energy

resources of natural systems. At the same time environmental assessment of landscapes productivity should be based on the use of geographical regularities manifested in the scale of territorial units of different hierarchical ranks, that is, in geo-ecological, environmental and landscape systems, which gives the possibility to explain the nature of the formation and functioning of landscape systems in specific climatic zones [1, 2].

Purpose of this study is to develop integrated models to assess biological and ecological productivity of landscapes, including plant productivity and soil conditions, allowing to use of qualitative and quantitative indicators of heat and moisture availability and to determine the regularities of formation and functioning of natural systems depending on the latitudinal zonation and altitudinal belts for the effective distribution of productive forces of agricultural sector.

Methods. Model development and assessment of biological and ecological capacity of natural system landscapes is based on the methodology of systemic researches in the field of biology, geography and ecology, as well as on the methods of mathematical modelling of natural process.

Long-term data of meteorological monthly bulletins at the meteorological stations of southern Kazakhstan, submitted to "Kazgidromet" RSE is used to assess the biological and ecological capacity of southern Kazakhstan natural systems landscapes [3, 4].

Results. Energy characteristics of the natural system are used for quantitative assessment of biological and ecological potential of landscapes, that is, the formation of the production process of the vegetative and soil covers in the landscape systems, that is: sum of air temperatures (Σt) above 10°C, the amount of moisture deficit of air (Σd , mb), evaporation (E_o , mm), photosynthetic active radiation (R , kJ/cm²) and the sum of precipitation (O_c , mm) (table 1) [4-10].

As can be seen from table 1, energy resources of South Kazakhstan landscape systems are very high, as the sum of biologically active temperatures ($\Sigma t, ^\circ C$) is in the range of 1737-4419°C, the sum of air humidity deficit (Σd) – 1190-4240 mb, evaporation (E_o) – 521-1325 mm and photosynthetic active radiation (R) - 115,8-204.6 kJ/cm², which have inverse dependences to the absolute height (H , m) of meteorological stations location. Precipitation (O_c - 151-509 mm) has a direct dependence, showing strict obedience to the laws of geographic zonality, which allows to use them for quantitative and qualitative assessment of biological and environmental productivity of vegetation and soil cover of southern Kazakhstan landscape systems.

Methodology for biological and environmental assessment of landscapes agricultural productivity was used for agroclimatic assessment of the landscapes productivity. This approach is used in mathematical models of plant and soil productivity of J. C. Mustafayev and G. A. Adilbektegi These models are based on the concept of maximum productivity. Similar studies conducted by scientists of CIS and foreign countries [11-20].

Biological productivity according to the official definition of the International coordinating committee on terminology and concepts in the field of production studies, is a set of processes of creation, transformation, absorption and transmission of energy through ecological and biological systems of different levels – from individual organisms to biogeocoenosis (ecosystems). The study of biological productivity of natural systems is a necessary basis of rational use, protection and reproduction of biological resources of the Earth.

To assess heat availability of landscape systems vegetation cover you can use the average annual amount of biologically active air temperatures of the $\sum t_i, ^\circ C$ region to the average maximum possible biologically active sums of air temperatures in the $\sum t_{\max}, ^\circ C$ region, that is: $C_{ts} = \sum t_i / \sum t_{\max}$. Expected productivity of vegetation, depending on the heat availability of landscape systems are determined by indicators (C_{tb}) characterizing favorable temperature regime of the natural system (1):

$$C_{tb} = 1 - C_{ts} = 1 - (\sum t_i / \sum t_{\max}), \quad (1)$$

To assess the moisture availability of the landscape systems soil cover you can use the natural moisturizing factor of N. N. Ivanov (2):

$$C_m = O_c / E_o, \quad (2)$$

where O_c is precipitation, mm; E_o - evaporation rate, determined by the formula of N.N. Ivanov (3) [8]:

Table 1 – Natural energy resources of Southern Kazakhstan landscape systems

Weather stations	Absolute height (H), m	Indicators of natural and energy resources				
		Σt , °C	Σd , mb	E_o , mm	R , kJ/cm ²	O_c , mm
Almaty region						
Uch-Aral	395	3294	2452	988	167,4	385
Sarkand	764	3163	2217	949	163,0	535
Taldykurgan	601	3173	2315	952	163,3	412
Bakanas	396	3525	2962	1058	175,0	223
Zharkent	641	3631	2396	1089	178,5	191
Chilik	606	3623	2559	1089	178,2	298
Almaty	671	3007	1370	902	157,9	509
Narynkol	1806	1737	1190	521	115,8	433
Sary-Ozek	548	2134	2257	640	129,0	332
Zhambyl region						
Ulanbel	266	3721	3050	1116	181,5	224
Moiynqum	350	3506	2553	1052	174,4	294
Uyuk	373	3720	3203	1116	181,5	283
Otar	742	3116	2635	935	161,5	316
Kurdai	1141	2930	2214	879	155,3	290
Kulan	682	3386	2519	1051	170,4	361
Taraz	642	3492	2309	1048	173,9	353
Merke	703	3472	2513	1041	173,2	435
Zhualy	952	2766	2022	830	149,9	447
Turkestan region						
Suzak	316	3822	3541	1147	184,8	186
Turkestan	206	4350	4197	1305	202,3	238
Tyulkubas	789	3876	3233	1163	186,6	951
Arys	237	4419	4240	1325	204,6	275
Shymkent	543	4065	3382	1219	192,9	582
Shardara	238	4397	4168	1391	203,9	264
Tolebi	455	3655	2605	1096	179,3	336
Kyzylorda region						
Saksaul	78	3647	3233	1094	179,0	152
Aral sea	62	3524	2633	1057	175,0	166
Qazaly	66	3647	2733	1094	179,0	178
Zhusaly	101	3809	3403	1142	184,4	165
Kyzylorda	128	3766	3160	1129	183,0	151
Shieli	152	3883	3154	1165	186,8	174
Ak-Kum	173	4253	3861	1276	199,1	204

$$E_o = 0.0018(25 + t)^2(100 - a), \quad (3)$$

where t - the average monthly air temperature, °C; a - average monthly relative humidity, %.

Thus, biological assessment of landscapes productivity (C_{bp}) is determined by the ratio of such averaged indicator values as plant (C_{tb}) and soil (C_m) productivity factors (4):

$$C_{bp} = C_{tb} \cdot C_m, \quad (4)$$

To determine the biological productivity of landscape systems in southern Kazakhstan and its agro-ecological regions the information-analytical materials presented in table 1(table 2) were used as potential indicators.

As can be seen from table 2, the qualitative and quantitative values of biological productivity of plant (C_{tb}) and soil (C_m) covers, as well as the biological productivity of landscapes (C_{bp}) is strictly subject to the laws of geographic zonality, that is directly dependent on the absolute height (H) location of landscape systems.

Table 2 – Evaluation of southern Kazakhstan
and its agro-ecological regions landscape systems biological productivity

Weather stations	Absolute height (H), m	Indicators of energy resources			C_m	C_{bp}
		Σt , °C	C_{ts}	C_{tb}		
Almaty region						
Uch-Aral	395	3294	0,74	0,26	0,19	0,050
Sarkand	764	3163	0,72	0,28	0,28	0,078
Taldykurgan	601	3173	0,72	0,28	0,22	0,062
Bakanas	396	3525	0,80	0,20	0,11	0,022
Zharkent	641	3631	0,82	0,18	0,13	0,023
Chilik	606	3623	0,82	0,18	0,10	0,018
Almaty	671	3007	0,68	0,32	0,35	0,112
Narynkol	1806	1737	0,39	0,61	0,65	0,396
Sary-Ozek	548	2134	0,48	0,52	0,23	0,120
Zhambyl region						
Ulanbel	266	3721	0,34	0,66	0,10	0,066
Moiynqum	350	3506	0,79	0,310	0,13	0,040
Uyuk	373	3720	0,84	0,16	0,12	0,019
Otar	742	3116	0,71	0,19	0,22	0,042
Kurdai	1141	2930	0,66	0,34	0,30	0,102
Kulan	682	3386	0,77	0,23	0,21	0,070
Taraz	642	3492	0,79	0,21	0,20	0,042
Merke	703	3472	0,79	0,21	0,29	0,061
Zhualy	952	2766	0,63	0,37	0,23	0,099
Turkestan region						
Suzak	316	3822	0,86	0,14	0,09	0,013
Turkestan	206	4350	0,98	0,02	0,08	0,002
Tyulkubas	789	3876	0,88	0,12	0,22	0,026
Arys	237	4419	0,99	0,01	0,08	0,001
Shymkent	543	4065	0,92	0,08	0,15	0,012
Shardara	238	4397	0,99	0,01	0,08	0,001
Tolebi	455	3655	0,83	0,17	0,17	0,029
Kyzylorda region						
Saksaul	78	3647	0,83	0,17	0,08	0,014
Aral sea	62	3524	0,80	0,20	0,16	0,034
Qazaly	66	3647	0,82	0,18	0,08	0,014
Zhusaly	101	3809	0,86	0,14	0,06	0,008
Kyzylorda	128	3766	0,85	0,15	0,07	0,011
Shieli	152	3883	0,88	0,12	0,06	0,007
Ak-Kum	173	4253	0,96	0,04	0,06	0,002

Ecological productivity of landscapes is closely connected with the flow of energy passing through a particular ecosystem, that is, falling to the trophic chain. Energy is accumulated as organic compounds, which provide continuous production of biomass (living matter). This is one of the fundamental processes of the biosphere.

Energy consumed for soil formation, defined by the formula of V.P. Volobuyev [6] characterizes the productivity of the soil landscapes to some extent (5):

$$Q_i = R \cdot \exp(-\alpha_o \cdot \bar{R}), \quad (5)$$

where Q_i - energy consumed for soil formation, kJ/cm²; α_o - coefficient taking into account the soil surface condition.

Table 3 – Evaluation of southern Kazakhstan and its agro-ecological regions landscape systems biological productivity

Weather stations	Absolute height (H), m	Energy costs on soil formation (Q , kJ/cm ²)			C_m	C_{bp}
		where \bar{R}_i	where $\bar{R} = 1.0$	C_p		
Almaty region						
Uch-Aral	395	60,9	104,6	0,58	0,19	0,11
Sarkand	764	92,2	101,9	0,90	0,28	0,25
Taldykurgan	601	77,9	101,9	0,76	0,22	0,17
Bakanas	396	40,0	109,4	0,37	0,11	0,04
Zharkent	641	56,5	108,8	0,52	0,13	0,07
Chilik	606	56,4	111,4	0,51	0,10	0,05
Almaty	671	51,5	98,7	0,52	0,35	0,21
Narynkol	1806	70,2	72,4	0,97	0,65	0,76
Sary-Ozek	548	62,1	80,6	0,77	0,23	0,18
Zhambyl region						
Ulanbel	266	39,7	113,4	0,35	0,10	0,04
Moiynqum	350	57,6	109,0	0,53	0,13	0,07
Uyuk	373	54,4	113,4	0,48	0,12	0,07
Otar	742	54,4	109,3	0,74	0,22	0,19
Kurdai	1141	71,9	97,1	0,53	0,30	0,16
Kulan	682	69,9	106,5	0,56	0,21	0,15
Taraz	642	68,8	108,7	0,53	0,20	0,14
Merke	703	82,6	108,6	0,76	0,29	0,26
Zhualy	952	79,8	93,7	0,85	0,23	0,20
Turkestan region						
Suzak	316	28,8	115,5	0,25	0,09	0,04
Turkestan	206	40,8	126,4	0,32	0,08	0,01
Tyulkubas	789	130,2	116,6	1,00	0,22	0,22
Arys	237	54,5	127,9	0,43	0,08	0,03
Shymkent	543	74,6	120,6	0,52	0,15	0,07
Shardara	238	47,8	127,3	0,38	0,08	0,03
Tolebi	455	66,0	112,0	0,59	0,17	0,11
Kyzylorda region						
Saksaul	78	19,6	111,8	0,18	0,08	0,01
Aral sea	62	24,4	109,4	0,22	0,16	0,04
Qazaly	66	27,0	111,9	0,24	0,08	0,02
Zhusaly	101	22,3	115,3	0,19	0,06	0,01
Kyzylorda	128	18,9	114,4	0,16	0,07	0,01
Shieli	152	25,0	116,8	0,21	0,06	0,01
Ak-Kum	173	31,9	114,4	0,26	0,06	0,01

In a natural system, the principle of energy balance of heat and moisture is observed in natural conditions, where the radiation index of dryness (\bar{R}) is equal to 1.0. Therefore, the limit in the range of 0.9-1.0 can be taken as a criterion of the radiative index of dryness (\bar{R}).

Then, the potential energy expended on soil-forming process (Q_n), providing the potential productivity of the soil can be determined by expression (6):

$$Q_n = R \cdot \exp(-0.9 \cdot \alpha_o) . \quad (6)$$

Consequently, the ratio of energy consumed for soil formation under natural conditions (Q_i) to the potential energy expended in soil-forming process (Q_n) represents the productivity of the landscapes soil cover, i.e. (7), [7]

$$C_p = Q_i / Q_n, \quad (7)$$

Thus the natural moisturizing factor of N. N. Ivanov [8] can be used to evaluate the productivity of vegetation landscape systems (8):

$$C_m = O_c / E_o . \quad (8)$$

Thus, the ecological productivity of landscapes (C_{ep}) is determined by the ratio of such averaged indicator values as an indicator of the soil surface productivity (C_p) and the natural moisturization coefficient, which characterizes the productivity of vegetation (C_m) (9):

$$C_{ep} = C_m \cdot C_p . \quad (9)$$

On the basis of information and analytical materials presented in table 1, which characterize energy resources and heat and moisture availability of southern Kazakhstan landscape systems, environmental productivity (C_{ep}) of the landscape of South of Kazakhstan and its agro-ecological regions (table 3) is defined.

As can be seen from table 3, the analysis of the plant (C_m) and soil (C_p) covers ecological productivity calculation results, as well as ecological productivity of landscapes (C_{bp}), indicates that their qualitative and quantitative values, and biological productivity of landscapes, strictly obey the laws of geographic zonality and decrease with decreasing altitude (H) of landscape systems location.

Thus, the developed model of landscapes biological and ecological productivity climate index allows, first, to give quantitative values of qualitative changes of habitats; secondly, to assess the modelling the transformation of natural systems under climate change; third, to conduct landscape-ecological zoning of the natural systems that can be used to assess the biological and ecological productivity of landscape systems, allowing the rational location of the agriculture productive forces.

Discussion. For the territory of South Kazakhstan, spanning from the mountain foothills to arid desert zones, the most important soil-ecological indices are set, i.e. mathematical models of biological and ecological productivity of landscapes that define the resource potential of the area and productivity of soil and vegetation cover. System for assessing the biological and ecological productivity of landscapes has attempted to carry out a comprehensive accounting of climate, soil and plants which would more fully describe the environment the agricultural sector is operating in.

Г. Ә. Әділбектері¹, Ж. С. Мұстафаев², Т. К. Уватаева¹, Ж. Н. Дулатбекова¹, Jozef Mosiej³

¹РМК ШЖҚ «Л. Н. Гумилев атындағы Еуразия ұлттық университеті», Астана, Қазақстан,

²«Қазақ ұлттық аграрлық университеті» коммерциялық емес акционерлік қоғамы, Алматы, Қазақстан,

³Жаратылыстану ғылымдарының Варшава университеті, Варшава, Польша

ОҢТҮСТИК ҚАЗАҚСТАН ЛАНДШАФТТАРЫНЫҢ БИОЛОГИЯЛЫҚ ЖӘНЕ ЭКОЛОГИЯЛЫҚ ӘЛЕУЕТІН САНДЫҚ ЖӘНЕ САПАЛЫҚ БАҒАЛАУ

Аннотация. Жұмыста жоғары тиімді агроландшафттық жүйелерді жобалау және құрастыру кезінде табиғи корларды неғұрлым толық және тиімді пайдалану мақсатында Оңтүстік Қазақстанның топырақ-климаттық жағдайларындағы ландшафтық жүйелердің биологиялық және экологиялық әлеуетін бағалау

жүйесінің әдістемелік тәсілдері қарастырылды. Жылумен, жарықпен және ылғалмен қамтамасыз етудің сапалық және сандық сипаттамалық белгілерінің негізінде, топырақ және өсімдік жамылғысын қамтитын ландшафттық жүйелердің биологиялық және экологиялық өнімділігін бағалауға арналған математикалық үлгі әзірленді.

Ландшафттардың биологиялық өнімділігін бағалауға әзірленген үлгі ($K_{\text{б}}$) ландшафттық жүйелердің топырақ жамылғысының жылумен қамтамасыз етілуін ($K_{\text{тб}}$), өсімдік жамылғысының табиги ылғалдану дәрежесінің өлшемдік көрсеткішін (K_y), соңдай-ақ ландшафттардың экологиялық өнімділігін ($K_{\text{бн}}$) ескереді. Бұл үлгі топыракты түзуші жүргігे шығын болған қуатына (Q_n) және өсімдік жамылғысының табиги ылғалдану көрсеткішіне (K_y) негізделеді және табиги жүйенің табиги-әлеуетті қорларын бағалау үшін география және экология саласында кеңінен пайдаланылатын табиги жүйенің климаттық өнімділігінің сапалық және сандық үлгілерінің бірі болып табылады.

Ландшафттық жүйелердің биологиялық және экологиялық өнімділігін бағалауға арналып әзірленген үлгінің негізінде, құрамына Алматы, Жамбыл, Түркістан және Қызылорда облыстары кіретін Қазақстанның Онтүстік аймағында орналасқан 32 метеорологиялық бекеттердің көпжылдық деректік мәліметтерін пайдалану арқылы, оның табиги-корлық әлеуетін жан-жақты бағалау жүргізілді.

Ландшафттық жүйелердің биологиялық және экологиялық өнімділігін бағалаудың жетілдірілген әдістерін пайдалану агроэкологиялық аудандар бойынша ландшафттық жүйелердің биологиялық корларын негұрлым дәл анықтауға және табиги жүйенің табиги-корларының әлеуетін пайдаланудың тиімділігін бағалауға мүмкіндік береді.

Түйін сөздер: табигат, ландшафт, климат, өнімділік, биология, экология, потенциал, индекс.

Г. А. Адильбектегі¹, Ж. С. Мустафаев², Т. К. Уватаева¹, Ж. Н. Дулатбекова¹, Jozef Mosiej³

¹РГП на ПХВ «Евразийский национальный университет им. Л. Н. Гумилева», Астана, Казахстан

²НАО Казахский национальный аграрный университет, Алматы, Казахстан,

³Варшавский университет Естественных наук, Варшава, Польша

КОЛИЧЕСТВЕННАЯ И КАЧЕСТВЕННАЯ ОЦЕНКА БИОЛОГИЧЕСКОГО И ЭКОЛОГИЧЕСКОГО ПОТЕНЦИАЛА ЛАНДШАФТОВ ЮЖНОГО КАЗАХСТАНА

Аннотация. В работе рассматриваются методические подходы к системе оценки биологического и экологического потенциала ландшафтных систем в почвенно-климатических условиях Южного Казахстана с целью наиболее полного и эффективного использования естественно-природных ресурсов при проектировании и конструировании высокоеффективных агроландшафтных систем. На основе качественных и количественных индикаторов тепло-, свето- и влагообеспеченности разработана математическая модель для оценки биологической и экологической продуктивности ландшафтных систем, включающая продуктивности почвенного и растительного покровов.

При этом следует отметить, что разработанная модель биологической оценки продуктивности ландшафттов ($K_{\text{б}}$) учитывает теплообеспеченность почвенного покрова ландшафтных систем ($K_{\text{тб}}$), коэффициент естественного увлажнения растительного покровов (K_y), а также экологическую продуктивность ландшафттов ($K_{\text{бн}}$). Эта модель базируется на потенциально возможной энергии, затраченной на почвообразовательный процесс (Q_n) и коэффициент естественного увлажнения растительного покровов (K_y), является одним из модификаций качественных и количественных моделей климатической продуктивности природной системы, широко используемых в области географии и экологии для оценки природно-потенциальных ресурсов природной системы.

На основе разработанной модели биологической и экологической продуктивности ландшафтных систем выполнена всесторонняя оценка природно-ресурсного потенциала Южно-Казахстанской области, включающая Алматинскую, Жамбылскую, Туркестанскую и Кызылординскую области с использованием многолетних данных 32 метеорологических станции, расположенных на территории региона.

Использование усовершенствованных методов оценки биологической и экологической продуктивности ландшафтных систем позволяет более точно определить биологические ресурсы ландшафтных систем по агроэкологическим районам и оценить эффективность использования природно-ресурсного потенциала природной системы.

Ключевые слова: природа, ландшафт, климат, продуктивность, биология, экология, потенциал, индекс.

Information about authors:

Adilbektegi Gulmira Adilbekkizi, Candidate of Geographical Sciences, acting. Associate Professor of the L. N. Gumilyov Eurasian National University, Astana, Kazakhstan; adilbekova.gulmira@yandex.kz; <https://orcid.org/0000-0002-1521-0145>

Mustafaev Zhumakhan Suleimenovich, doctor of technical sciences of the Russian Federation and the RK, professor of the Kazakh National Agrarian University, Almaty, Kazakhstan; z-mustafa@rambler.ru; <https://orcid.org/0000-0003-2425-8148>

Uvatayeva Togzhan Kairatovna, Master of the 2nd course of the specialty 6M060800 - Ecology of the L. N. Gumilyov Eurasian National University, Astana, Kazakhstan; t_uvatayeva@mail.ru; <https://orcid.org/0000-0002-2404-6260>

Dulatbekova Zhanna Nurzhankuzy, Master of the 2nd course of the specialty 6M060800 - Ecology of the L. N. Gumilyov Eurasian National University, Astana, Kazakhstan; yadulatbekova@gmail.com; <https://orcid.org/0000-0002-2522-0822>

Mosiej Jozef, Professor of the Warsaw University of Natural Sciences, Warsaw, Poland; jozef_mosiej@sggw.pl; <https://orcid.org/0000-0002-8040-7032>

REFERENCES

- [1] Mustafayev Z.S., Adilbektegi G.A., Ryabcev A.D. (2007) Methodological basis for assessing the stability and resistance of landscapes. Taraz. ISBN: 9965-724-18-0.
- [2] Mustafayev Z.S., Ryabcev A.D., Kozykeeva A.T., Adilbektegi G.A. (2007) Landscapes and natural and man-made complexes. Taraz. ISBN: 9965-724-19-9.
- [3] Scientific and Applied Reference Book on Climate of the USSR (1989). Series 13, Long-term data. Part 1-6, issue. Book 2. Gidrometeoizdat, Russia (in Russ.).
- [4] Mustafayev Z.S. (2017) Melioration of agricultural land in Kazakhstan [Environmental Engineering] 1: 87-93 (in Russ.).
- [5] Shashko D.I. (1985) To take into account the bioclimatic potential. [Agriculture] 4: 19-26 (in Russ.).
- [6] Volobuyev V.R. (1974) Introduction to the energy of soil formation. M.: Nauka (in Russ.).
- [7] Buscha M., La Notte A., Laporte V., Erhard M. (2012) Potentials of quantitative and qualitative approaches to assessing ecosystem services // Ecological Indicators Journal. 21, 89-103. DOI: 10.1016/j.ecolind.2011.11.010 (in Eng.).
- [8] Ivanov N.N. (1941) Zones of humidification of the globe // Proceedings of the USSR Academy of Sciences. 3: 15-32 (in Russ.).
- [9] Hargrove W.W., Hoffman F.M. (2005). Potential of multivariate quantitative methods for delineation and visualization of ecoregions // Environ. Manag. 34, 39-60. DOI: 10.1007/s00267-003-1084-0 (in Eng.).
- [10] Long J., Nelson T., Wulder M. (2010). Regionalization of landscape pattern indices using multivariate cluster analysis // Environm. Manag. 46, 134-142. DOI: 10.1007/s00267-010-9510-6 (in Eng.).
- [11] Xu Ch., Sheng S., Chi T., Yang X., An S., Liu M. (2014). Developing a quantitative landscape regionalization framework integrating driving factors and response attributes of landscapes // Landscape Ecology Engineering. 10, 295-307. DOI: 10.1007/BF00135075 (in Eng.).
- [12] Raudsepp-Hearne C., Peterson G.D., Bennet E.M. (2010) Ecosystem service bundles for analyzing trade-offs in diverse landscapes // Proceedings of the National Academy of Sciences of the United States of America. 107, 5242-5247. DOI: 10.1073/pnas.0907284107 (in Eng.).
- [13] Perko D., Hrvatin M., Ciglic R. (2015) A methodology for natural landscape typification of Slovenia // Acta geographica Slovenica. 55-2. DOI: 10.3986/AGS.1938 (in Eng.).
- [14] Blum WEH Functions of soil for society and the environment (2005) Functions of soil for society and the environment // Rev Environ Sci Biotechnol. 4(3):75-79. DOI: 10.1007/s11157-005-2236-x (in Eng.).
- [15] Van Etvelde V., Antrop M. (2009) A stepwise multi-scaled landscape typology and characterization for transregional integration, applied on the federal state of Belgium // Landsc. Urban Plann. 91, 160-170. DOI: 10.1016/j.landurbplan.2008.12.008 (in Eng.).
- [16] Bunce R.G.H., Barr C.J., Clarke R.T., Howard D.C., Lane A.M.J. (1996) Land classification for strategic ecological survey // Journal of environmental management. 47-1. DOI: 10.1006/jema.1996.0034 (in Eng.).
- [17] Kwiatkowska-Malina J. (2018) Qualitative and quantitative soil organic matter estimation for sustainable soil management // Journal of Soils and Sediments. 18: 2801-2812. DOI: 10.1007/s11368-017-1891-1 (in Eng.).
- [18] Coleman D.C., Grys-Rubenstein E. (2014) Biological productivity, AccessScience. DOI:10.1036/1097-8542.082800 (in Eng.).
- [19] Tilman D., Reich P.B., Forest I. (2012) Biodiversity impacts ecosystem productivity as much as resources, disturbance, or herbivory // Proc Natl Acad Sci U S A 109(26): 10394-10397. DOI: 10.1073/pnas.1208240109 (in Eng.).
- [20] Bünenmann E.K., Bongiorno G., Zhanguo B., Brussaard L. (2018) Soil quality – A critical review // Soil Biology and Biochemistry. 120:105-125. DOI: 10.1016/j.soilbio.2018.01.030 (in Eng.).

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka-nanrk.kz

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

<http://www.geolog-technical.kz/index.php/en/>

Верстка Д. Н. Калкабековой

Подписано в печать 15.11.2019.
Формат 70x881/8. Бумага офсетная. Печать – ризограф.
19,7 п.л. Тираж 300. Заказ 6.